CHATGPT

Ankylosing spondylitis pain

Ankylosing spondylitis (AS) is a type of inflammatory arthritis that primarily affects the spine, causing pain and stiffness. Here are some common aspects related to the pain:

1. **Nature of Pain**: The pain often starts in the lower back and buttocks and can spread to the hips and shoulders. It's usually worse at night or in the early morning and can improve with movement.
2. **Duration**: The pain can vary in intensity and may last for hours or even days, often improving with activity.
3. **Inflammation**: AS causes inflammation in the spinal joints, which can lead to reduced flexibility and, over time, possible fusion of the spine.
4. **Management**: Treatment options typically include physical therapy, anti-inflammatory medications, and sometimes biologics or DMARDs to help reduce inflammation and manage pain.
5. **Lifestyle Changes**: Regular exercise and good posture can be beneficial. Heat and cold therapy may also help alleviate symptoms.

If you or someone you know is experiencing symptoms of AS, it's important to consult a healthcare professional for an accurate diagnosis and appropriate management.

Ankylosing spondylitis symptoms

Ankylosing spondylitis (AS) is a type of inflammatory arthritis primarily affecting the spine and sacroiliac joints. Its hallmark feature is chronic inflammation, which can lead to the fusion of vertebrae (ankylosis), causing a loss of spinal mobility. Common symptoms of AS include:

1. **Back Pain and Stiffness**: This is typically worse in the morning or after periods of inactivity. The pain often starts in the lower back and buttocks and may gradually spread up the spine.
2. **Improvement with Exercise**: Unlike mechanical back pain, AS pain tends to improve with movement and exercise.
3. **Reduced Spinal Mobility**: Over time, the inflammation can cause the vertebrae to fuse, leading to a loss of flexibility in the spine, a condition known as "bamboo spine."
4. **Pain in Other Joints**: AS can also affect peripheral joints such as the hips, shoulders, knees, and ribs, causing pain and swelling.
5. **Fatigue**: Chronic inflammation and pain can lead to a general sense of fatigue.
6. **Neck Pain**: In some cases, neck stiffness and pain may also be present.

7. **Limited Chest Expansion**: Inflammation in the joints between the ribs and the spine can limit chest expansion, leading to difficulty in deep breathing.
8. **Eye Inflammation (Uveitis/Iritis)**: AS is commonly associated with inflammation in the eye, causing redness, pain, light sensitivity, and blurred vision.
9. **Enthesitis**: Inflammation at the sites where tendons or ligaments attach to the bone, commonly seen in the heel (Achilles tendon) or bottom of the foot (plantar fascia).
10. **Postural Changes**: Severe cases of AS can lead to a forward stooped posture due to spinal fusion.

If left untreated, AS can result in significant limitations in daily activities and a reduced quality of life due to the progressive stiffening of the spine. Early diagnosis and treatment are essential for managing symptoms and slowing disease progression.

Ankylosing spondylitis disease

Ankylosing spondylitis (AS) is a type of inflammatory arthritis that primarily affects the spine and the sacroiliac joints in the pelvis. It can cause pain and stiffness in the back and neck, particularly in the early morning or after periods of inactivity. Over time, AS can lead to the fusion of the vertebrae, reducing flexibility and potentially resulting in a hunched posture.

Key Features:

- **Symptoms**: Chronic back pain, stiffness, fatigue, and sometimes peripheral joint pain. Symptoms can improve with physical activity but worsen with rest.
- **Causes**: The exact cause is unknown, but genetics plays a significant role. A majority of people with AS have the HLA-B27 gene, although not everyone with the gene will develop the condition.
- **Diagnosis**: Diagnosis typically involves a combination of medical history, physical examinations, imaging studies like X-rays or MRIs, and blood tests for inflammatory markers and the HLA-B27 antigen.
- **Treatment**: While there's no cure, treatments aim to reduce pain and stiffness. Common approaches include nonsteroidal anti-inflammatory drugs (NSAIDs), physical therapy, and, in some cases, biologic medications that target specific components of the immune system.

Living with AS:

Managing AS often involves lifestyle changes, including regular exercise, maintaining good posture, and sometimes using assistive devices.

Support groups and education can also be beneficial for those living with the condition.

If you have specific questions about symptoms, treatment options, or managing AS, feel free to ask!

What is Ankylosing spondylitis

Ankylosing spondylitis (AS) is a chronic inflammatory disease that primarily affects the spine and sacroiliac joints, causing pain and stiffness. It is a type of arthritis, specifically classified as a **seronegative spondyloarthropathy**, meaning it does not involve the presence of rheumatoid factor in the blood.

Key Features of Ankylosing Spondylitis:

- **Inflammation of the spine**: The disease leads to inflammation in the joints between the vertebrae (spinal bones), which can result in the fusion of these bones over time, leading to reduced flexibility and a hunched posture.
- **Sacrolitis**: Early signs often involve inflammation in the sacroiliac joints (where the spine meets the pelvis), causing lower back pain.
- **Enthesitis**: Inflammation at the sites where tendons and ligaments attach to bones, particularly in areas like the Achilles tendon or plantar fascia.
- **Pain and stiffness**: AS typically causes chronic back pain and morning stiffness that improves with exercise. Unlike mechanical back pain, it often worsens with rest.
- **Progressive spinal fusion**: Over time, ankylosing spondylitis can cause the spine to become less flexible, and in advanced cases, the vertebrae can fuse together (ankylosis), leading to a loss of mobility.

Risk Factors and Causes:

- **Genetic predisposition**: The HLA-B27 gene is strongly associated with AS. However, not everyone with the gene develops the disease.
- **Gender and age**: AS is more common in men and usually starts in early adulthood (late teens to 30s).

Symptoms:

- **Chronic back pain**, especially in the lower back and buttocks.
- **Morning stiffness** lasting more than 30 minutes.
- **Fatigue** and possible weight loss in advanced stages.
- **Reduced chest expansion** due to involvement of the ribs.
- **Other joint involvement**: In some cases, it can also affect peripheral joints like the hips, shoulders, or knees.

Diagnosis:

- **Imaging**: X-rays or MRI can show characteristic changes in the sacroiliac joints and spine, such as sacroiliitis or "bamboo spine" (fusion of vertebrae).
- **Blood tests**: Testing for HLA-B27, elevated inflammatory markers (CRP, ESR).

Treatment:

- **Non-steroidal anti-inflammatory drugs (NSAIDs)**: The first line of treatment to reduce pain and inflammation.
- **Physical therapy**: To maintain mobility and posture.
- **TNF inhibitors and biologics**: For more severe cases, these medications can reduce inflammation and slow disease progression.
- **Surgery**: Rarely, in advanced cases with severe joint damage or spinal deformity.

Ankylosing spondylitis is a lifelong condition, but with proper management, patients can lead a relatively normal life with improved symptoms.

Ankylosing spondylitis treatment

Ankylosing spondylitis (AS) is a chronic inflammatory disease primarily affecting the spine and sacroiliac joints, leading to pain and stiffness. The treatment aims to relieve symptoms, reduce inflammation, prevent deformities, and maintain functional mobility. Key components of AS treatment include:

1. Medications

- **Non-Steroidal Anti-Inflammatory Drugs (NSAIDs):** First-line treatment to reduce pain and inflammation. Examples include ibuprofen, naproxen, and indomethacin.
- **Tumor Necrosis Factor (TNF) Inhibitors:** Biologic drugs like etanercept, infliximab, and adalimumab are prescribed when NSAIDs are ineffective.
- **Interleukin-17 (IL-17) Inhibitors:** Secukinumab and ixekizumab are newer biologics targeting IL-17, a key player in AS inflammation.
- **Disease-Modifying Anti-Rheumatic Drugs (DMARDs):** Used in peripheral joint involvement; sulfasalazine and methotrexate are commonly used.
- **Corticosteroids:** Short-term use for severe flare-ups or localized joint inflammation, although long-term use is avoided due to side effects.

2. Physical Therapy and Exercise

- **Stretching and Strengthening Exercises:** To maintain flexibility and strength, particularly in the spine and large joints.
- **Postural Training:** Exercises that promote good posture help to reduce spinal deformity progression.
- **Aerobic Exercise:** Swimming, cycling, or walking can improve overall fitness without stressing the spine.

3. Lifestyle Modifications

- **Quit Smoking:** Smoking can worsen the progression of ankylosing spondylitis.
- **Healthy Diet:** Anti-inflammatory foods may help reduce overall inflammation.
- **Ergonomic Adjustments:** Modifications in sleeping and sitting positions can help reduce discomfort.

4. Surgical Intervention

- **Joint Replacement:** In cases where the hip or knee joints are severely damaged, joint replacement surgery may be needed.
- **Spinal Surgery:** Rarely, surgery may be needed for spinal deformity correction, such as severe kyphosis, or if neurological symptoms develop due to nerve compression.

5. Patient Education and Support

- Education on disease management, support groups, and counseling can help patients cope with the chronic nature of the condition.

Treatment is individualized based on disease severity, patient response, and functional limitations.

As

It seems like your message was cut off. Could you please clarify what you were going to ask?

As Ankylosing spondylitis

Ankylosing spondylitis (AS) is a type of inflammatory arthritis that primarily affects the spine, leading to pain and stiffness. It can also affect other joints and areas, including the hips, shoulders, and sometimes even organs like the eyes.

Symptoms:

- Chronic pain and stiffness in the lower back and hips, especially in the morning or after periods of inactivity
- Improved symptoms with movement

- Fatigue
- Reduced flexibility in the spine over time
- In some cases, inflammation of the eyes (iritis or uveitis)

Causes:

The exact cause of AS isn't fully understood, but it's believed to involve a combination of genetic and environmental factors. A significant number of people with AS carry the HLA-B27 antigen.

Management:

While there's no cure, treatment options can help manage symptoms:

- **Medications**: Nonsteroidal anti-inflammatory drugs (NSAIDs) are commonly used, and biologics may be prescribed for more severe cases.
- **Physical therapy**: Regular exercise can improve flexibility and reduce pain.
- **Lifestyle changes**: Maintaining a healthy weight and staying active can help manage symptoms.

If you or someone you know is dealing with AS, it's important to work closely with a healthcare provider to create a personalized management plan. Would you like to know more about a specific aspect of AS?

Ankylosing spondylitis meaning

Ankylosing spondylitis is a type of inflammatory arthritis that primarily affects the spine, leading to pain and stiffness. Over time, it can cause some of the vertebrae in the spine to fuse together, resulting in reduced flexibility and a hunched posture. It often starts in early adulthood and can also affect other joints and areas, such as the hips and shoulders. The exact cause is not fully understood, but genetics may play a role. Treatment typically focuses on managing symptoms and maintaining mobility.

Ankylosing spondylitis causes

The exact cause of ankylosing spondylitis isn't fully understood, but it involves a combination of genetic predisposition (especially the HLA-B27 gene), immune system dysfunction, and potential environmental triggers. If you're interested, I can delve into how these factors interact.

Ankylosing spondylitis diagnosis

Ankylosing spondylitis (AS) is diagnosed through a combination of clinical assessment, imaging studies, and laboratory tests. Here's a breakdown of the diagnostic process:

1. Medical History

- **Symptoms**: Patients often report chronic back pain, stiffness (especially in the morning), and pain that improves with exercise.
- **Family History**: A family history of AS or related conditions can be significant.

2. Physical Examination

- **Range of Motion**: Assessment of spinal flexibility and posture.
- **Tenderness**: Checking for tenderness in the spine and sacroiliac joints.

3. Imaging Studies

- **X-rays**: Initial imaging can show changes in the sacroiliac joints, which may indicate inflammation.
- **MRI**: More sensitive than X-rays, MRIs can detect early inflammatory changes before structural damage occurs.

4. Laboratory Tests

- **HLA-B27 Testing**: Presence of the HLA-B27 antigen is associated with AS, but not all patients with AS test positive for it.
- **Inflammatory Markers**: Blood tests for C-reactive protein (CRP) or erythrocyte sedimentation rate (ESR) can indicate inflammation.

5. Diagnostic Criteria

- The **Modified New York Criteria** are often used, which include:
  - Low back pain and stiffness for more than three months.
  - Improvement with exercise but not with rest.
  - Limited range of motion in the lumbar spine.
  - Limited chest expansion.

Summary

A comprehensive approach combining these elements helps confirm the diagnosis of ankylosing spondylitis. Early diagnosis and treatment are important to manage symptoms and prevent complications. If you have any specific concerns or need more detailed information, feel free to ask!

Ankylosing spondylitis icd10

The ICD-10 code for ankylosing spondylitis is **M45**. This category includes various subtypes, such as:

- **M45.0**: Ankylosing spondylitis with involvement of the cervical spine
- **M45.1**: Ankylosing spondylitis with involvement of the thoracic spine

- **M45.2**: Ankylosing spondylitis with involvement of the lumbar spine
- **M45.9**: Ankylosing spondylitis, unspecified

If you need more specific details or assistance, feel free to ask!

Ankylosing spondylosis

Ankylosing spondylosis, commonly known as ankylosing spondylitis, is a type of inflammatory arthritis that primarily affects the spine. Here’s an overview:

Symptoms

- **Chronic Pain and Stiffness**: Typically starts in the lower back and can improve with movement.
- **Reduced Flexibility**: Over time, the spine may become less flexible, leading to a hunched posture.
- **Fatigue**: Chronic inflammation can cause general fatigue.
- **Possible Joint Involvement**: Other joints, such as hips and shoulders, may also be affected.

Causes

- **Genetics**: A strong association with the HLA-B27 gene.
- **Immune System**: The body’s immune response may play a role in the development of inflammation.

Diagnosis

- **Medical History and Physical Exam**: Assessment of symptoms and physical limitations.
- **Imaging Tests**: X-rays or MRIs can reveal changes in the spine and pelvis.
- **Blood Tests**: To check for markers like HLA-B27.

Treatment

- **Medications**: Nonsteroidal anti-inflammatory drugs (NSAIDs) are commonly used. Disease-modifying antirheumatic drugs (DMARDs) and biologics may also be prescribed for more severe cases.
- **Physical Therapy**: Exercises to improve flexibility and posture.
- **Lifestyle Changes**: Regular exercise, maintaining a healthy weight, and avoiding smoking can help manage symptoms.

Management

- Regular follow-ups with a healthcare provider.
- Monitoring for potential complications, such as osteoporosis or eye inflammation.

If you have specific questions or need more details about any aspect, feel free to ask!

Hla b27

HLA-B27 is a protein that can be associated with certain autoimmune diseases, particularly ankylosing spondylitis and other spondyloarthropathies. Would you like to delve deeper into its significance or related conditions?

Uveitis

Uveitis is an inflammation of the uvea, the middle layer of the eye. It can affect various parts of the uvea, including the iris (iritis), ciliary body (cyclitis), and choroid (choroiditis). Here are some key points about uveitis:

Symptoms

- **Eye redness**
- **Pain** or discomfort
- **Light sensitivity** (photophobia)
- **Blurred vision**
- **Floaters** (spots in your vision)

Causes

Uveitis can be caused by:

- **Infections** (viral, bacterial, or fungal)
- **Autoimmune diseases** (like rheumatoid arthritis or lupus)
- **Trauma** to the eye
- **Certain medications**

Diagnosis

Diagnosis typically involves:

- A comprehensive eye exam
- Medical history review
- Imaging tests (like optical coherence tomography)

Treatment

Treatment depends on the underlying cause and severity and may include:

- **Corticosteroids** (eye drops, oral medications, or injections)
- **Immunosuppressive drugs** for autoimmune conditions

- **Antibiotics** or antiviral medications if an infection is present
- **Surgery** in severe cases or if complications arise

Prognosis

With prompt treatment, many people recover well, but complications like glaucoma, cataracts, or vision loss can occur if not managed properly.

If you have specific questions or need more detailed information, feel free to ask!

Ankylosing spondylitis radiology

Ankylosing spondylitis (AS) is primarily evaluated through radiological imaging, which helps in diagnosing and monitoring the disease. Here are some key points regarding radiology in AS:

Imaging Techniques

1. **X-rays**:
   - **Initial Assessment**: X-rays are commonly used for initial evaluation.
   - **Typical Findings**:
     - Sacroiliitis: Inflammation of the sacroiliac joints is a hallmark feature.
     - Bamboo spine: Advanced cases may show fusion of the vertebrae, giving a characteristic "bamboo" appearance.
     - Syndesmophytes: Vertical bony growths can appear between vertebrae.
2. **MRI**:
   - **Sensitivity**: MRI is more sensitive than X-rays for early detection of inflammation.
   - **Findings**:
     - Bone marrow edema in the sacroiliac joints indicates active inflammation.
     - Early changes may include enthesitis (inflammation at tendon insertions).
3. **CT Scans**:
   - **Detailed Evaluation**: CT can provide detailed images of the spine and sacroiliac joints, especially for assessing structural changes.

Common Radiological Findings

- **Sacroiliac Joint Changes**:
  - Early: Blurring of joint margins, subchondral sclerosis.
  - Late: Joint fusion and irregularity.
- **Spinal Changes**:
  - Anterior vertebral corner involvement.

    - Progression leads to syndesmophytes and eventual fusion of the spine.
- **Extraskeletal Manifestations**:
    - May include findings in the chest (costovertebral joint involvement) and peripheral joints.

Interpretation

- Radiological findings are assessed in conjunction with clinical symptoms and lab tests (like HLA-B27).
- Radiologists play a crucial role in recognizing and interpreting these changes, which can guide treatment and management.

If you need more detailed information on a specific aspect, feel free to ask!

Spondyloarthritis

Spondyloarthritis refers to a group of inflammatory rheumatic diseases that primarily affect the spine and joints. Here are some key points:

Types

1. **Ankylosing Spondylitis (AS)**: The most common form, primarily affecting the spine.
2. **Psoriatic Arthritis**: Associated with psoriasis and can affect the joints and spine.
3. **Reactive Arthritis**: Follows an infection, often in the urogenital or gastrointestinal tract.
4. **Enteropathic Arthritis**: Linked to inflammatory bowel diseases like Crohn's disease and ulcerative colitis.

Symptoms

- **Chronic Pain**: Especially in the lower back and hips.
- **Stiffness**: Often worse in the morning or after periods of inactivity.
- **Reduced Flexibility**: Particularly in the spine.
- **Fatigue**: Common among those affected.
- **Peripheral Joint Pain**: In some forms, like psoriatic arthritis.

Diagnosis

Diagnosis usually involves:

- **Medical History**: Symptoms and family history.
- **Physical Exam**: Checking for spinal mobility and joint tenderness.
- **Imaging**: X-rays or MRIs to assess changes in the spine and joints.
- **Blood Tests**: Looking for specific markers like HLA-B27.

Treatment

Treatment aims to reduce inflammation and relieve pain:

- **Medications**: NSAIDs, disease-modifying antirheumatic drugs (DMARDs), and biologics.
- **Physical Therapy**: To maintain flexibility and strength.
- **Lifestyle Changes**: Exercise, maintaining a healthy weight, and smoking cessation can help manage symptoms.

Prognosis

With early diagnosis and appropriate treatment, many people can manage symptoms effectively and maintain a good quality of life.

If you have specific questions or need more detailed information, feel free to ask!

Ankylosing spondylitis blood test

Ankylosing spondylitis (AS) is an inflammatory arthritis primarily affecting the spine and sacroiliac joints. A key blood test associated with AS is the HLA-B27 test. While having this marker can increase the likelihood of developing AS, not everyone with AS tests positive for HLA-B27, and not everyone who tests positive will develop the condition.

Other blood tests may include:

1. **C-reactive protein (CRP)**: Measures inflammation in the body.
2. **Erythrocyte sedimentation rate (ESR)**: Another indicator of inflammation.

These tests help assess inflammatory activity but are not definitive for diagnosing AS. Diagnosis typically involves a combination of medical history, physical examination, imaging studies (like X-rays or MRIs), and sometimes genetic testing. If you suspect you have AS, it’s best to consult a healthcare professional for a thorough evaluation.

GEMİNİ

Ankylosing spondylitis pain +

Ankylosing Spondylitis Pain: Understanding and Managing

Ankylosing spondylitis (AS) is a chronic inflammatory condition primarily affecting the spine. It can cause significant pain and discomfort.

Common Pain Patterns

Lower back pain: This is often the first symptom and can be severe, especially in the morning or after periods of inactivity.

Stiffness: Stiffness in the spine, particularly in the lower back and hips, is common, especially upon waking.

Neck pain: As the condition progresses, neck pain may also develop.

Joint pain: Pain can also occur in other joints, such as the hips, shoulders, and knees.

Factors Affecting Pain

Flare-ups: Periods of increased pain and inflammation are common.

Stress: Emotional stress can exacerbate symptoms.

Weather: Changes in weather, particularly cold and damp conditions, may trigger pain.

Activity level: Excessive physical activity or prolonged periods of inactivity can contribute to pain.

Managing Ankylosing Spondylitis Pain

While there's no cure for AS, various strategies can help manage pain and improve quality of life:

Medication: Nonsteroidal anti-inflammatory drugs (NSAIDs), disease-modifying antirheumatic drugs (DMARDs), and biologic agents can help reduce inflammation and pain.

Physical therapy: Exercises can improve flexibility, strengthen muscles, and improve posture.

Occupational therapy: This can help individuals adapt their daily activities to minimize pain.

Hot and cold therapy: Applying heat or cold to the affected areas can provide temporary relief.

Lifestyle modifications: Maintaining a healthy weight, getting enough sleep, and managing stress can also help.

If you're experiencing symptoms of AS, it's important to consult with a healthcare professional for a proper diagnosis and treatment plan.

Would you like to learn more about specific treatment options or lifestyle changes that can help manage AS pain?

Ankylosing spondylitis symptoms

Ankylosing spondylitis is a chronic inflammatory condition that affects the spine and other joints. It can cause pain, stiffness, and fatigue. Some of the common symptoms include:

Back pain and stiffness: This is often the first symptom to appear. The pain is usually worse in the morning and after periods of inactivity.

Fatigue: Feeling tired all the time is a common symptom of ankylosing spondylitis.

Reduced range of motion: As the condition progresses, it can become difficult to move the spine and other joints.

Stooped posture: The spine may become curved, leading to a stooped posture.

Eye inflammation: Uveitis, a type of eye inflammation, can occur in people with ankylosing spondylitis.

Heel pain: Pain in the heel of the foot is another possible symptom.

Heart problems: Ankylosing spondylitis can increase the risk of heart problems, such as aortic valve disease.

It's important to see a doctor if you have any of these symptoms. Early diagnosis and treatment can help manage the condition and prevent complications.

Ankylosing spondylitis disease +

Ankylosing Spondylitis: A Chronic Inflammatory Disease

Ankylosing spondylitis is a chronic inflammatory disease that primarily affects the spine. Over time, it can cause the vertebrae in the spine to fuse together, leading to stiffness and loss of flexibility.

Symptoms

Back pain and stiffness: This is often worse in the morning and improves with activity.

Fatigue: Feeling tired even with enough sleep.

Reduced range of motion: Difficulty turning the head or twisting the body.

Eye inflammation: Iritis or uveitis can cause eye pain, redness, and blurred vision.

Heel pain: Enthesitis, inflammation where a tendon attaches to a bone, can cause heel pain.

Causes

The exact cause of ankylosing spondylitis is unknown, but genetic factors play a significant role. It is more common in people with a genetic marker called HLA-B27.

Treatment

While there's no cure for ankylosing spondylitis, treatment can help manage symptoms and improve quality of life. Common treatments include:

Medications: Nonsteroidal anti-inflammatory drugs (NSAIDs), disease-modifying antirheumatic drugs (DMARDs), and biologic therapies.

Physical therapy: Exercises to maintain flexibility, improve posture, and strengthen muscles.

Occupational therapy: Strategies to help with daily activities and adapt to changes in mobility.

If you suspect you may have ankylosing spondylitis, it's important to see a healthcare professional for diagnosis and treatment. Early diagnosis and treatment can help prevent complications and improve your overall well-being.

Would you like to know more about specific treatment options or the impact of ankylosing spondylitis on daily life?

What is Ankylosing spondylitis +

Ankylosing spondylitis (AS) is a type of arthritis that primarily affects the spine. It causes inflammation of the joints in the spine, particularly where the spine meets the pelvis. Over time, this inflammation can lead to the bones in the spine fusing together, making the spine less flexible and causing a hunched posture.

Key symptoms of AS include:

Back pain and stiffness, especially in the morning

Limited range of motion in the spine

Fatigue

Eye inflammation (uveitis)

Digestive problems (e.g., diarrhea, constipation)

Skin rashes

While the exact cause of AS is unknown, it is believed to be related to genetics. People with a family history of AS are at a higher risk.

Treatment for AS focuses on managing symptoms and preventing complications. This may involve a combination of:

Medications (e.g., nonsteroidal anti-inflammatory drugs, disease-modifying anti-rheumatic drugs, biologic agents)

Physical therapy (to improve flexibility and strength)

Occupational therapy (to help with daily activities)

If you are experiencing symptoms of AS, it is important to see a doctor for diagnosis and treatment.

Ankylosing spondylitis treatment +

Ankylosing Spondylitis Treatment

Ankylosing spondylitis (AS) is a chronic autoimmune condition that primarily affects the spine. While there's no cure, various treatments can help manage symptoms, improve quality of life, and prevent complications.

Treatment Options

Medications:

Nonsteroidal Anti-Inflammatory Drugs (NSAIDs): These medications help reduce pain and inflammation.

Disease-Modifying Anti-Rheumatic Drugs (DMARDs): These medications can slow the progression of the disease.

Biologics: These are newer medications that target specific parts of the immune system to reduce inflammation.

Corticosteroids: These can be used for short periods to manage severe flare-ups.

Physical Therapy: Regular exercise and stretching can help maintain flexibility, improve posture, and strengthen muscles.

Occupational Therapy: This can help individuals with AS learn to adapt to daily activities and manage pain.

Surgery: In severe cases, surgery may be necessary to correct spinal deformities or replace damaged joints.

Lifestyle Modifications

Regular Exercise: Physical activity can help maintain flexibility, reduce pain, and improve overall well-being.

Proper Posture: Maintaining good posture can help prevent spinal deformities.

Heat Therapy: Applying heat to sore joints can help relieve pain and stiffness.

Weight Management: Maintaining a healthy weight can reduce stress on the joints.

It's important to consult with a healthcare professional for a personalized treatment plan. They can assess your specific situation and recommend the most appropriate options.

Would you like to know more about any of these treatment options or have other questions?

As

Here are some of the most common meanings of "as":

Conjunctions:

Like: "She sings as beautifully as a bird."

Because: "As it was raining, we stayed at home."

While: "As I was walking, I saw a dog."

When: "As soon as I arrived, I called you."

Although: "As I like chocolate, I don't eat it often."

Prepositions:

In the role of: "He works as a teacher."

At the same time as: "We arrived at the same time as them."

In the manner of: "She dances as gracefully as a swan."

Adverbs:

So: "As soon as possible."

Also: "We went to the cinema and as well as watching a film, we had dinner."

Other uses:

In comparisons: "The house is as big as a castle."

To introduce clauses: "As I said before,..."

In legal documents: "As per the contract..."

Examples:

"As I was walking down the street, I saw a cat."

"She is as tall as her mother."

"He works as a doctor."

"As soon as I finish my homework, I can go out."

"Although it was raining, we went for a walk."

I hope this helps!

As Ankylosing spondylitis +

Ankylosing Spondylitis: A Brief Overview

Ankylosing spondylitis (AS) is a chronic inflammatory disease that primarily affects the spine. It causes inflammation in the joints, leading to pain, stiffness, and potential loss of mobility.

Key Symptoms

Back pain: Often begins in the lower back and can worsen over time.

Stiffness: Especially noticeable in the morning or after periods of inactivity.

Fatigue: Feeling tired or exhausted.

Reduced range of motion: Difficulty turning the head or bending the back.

Eye inflammation: Iritis or uveitis can occur in some individuals.

Causes and Risk Factors

Genetics: A specific gene, HLA-B27, is often present in people with AS.

Inflammation: The exact cause of the inflammation is unknown, but it is believed to be related to an autoimmune response.

Treatment

While there's no cure for AS, treatment focuses on managing symptoms and improving quality of life. This may include:

Medications: Nonsteroidal anti-inflammatory drugs (NSAIDs), disease-modifying antirheumatic drugs (DMARDs), and biologic therapies.

Physical therapy: Exercises to maintain flexibility and strengthen muscles.

Occupational therapy: Strategies to help with daily activities.

Surgery: In severe cases, surgery may be necessary to correct spinal deformities or replace damaged joints.

If you suspect you may have AS, it's important to consult a healthcare professional for a proper diagnosis and treatment plan.

Would you like to know more about a specific aspect of AS, such as its diagnosis, treatment options, or daily living with the condition?

Ankylosing spondylitis meaning

Ankylosing spondylitis (AS) is a type of arthritis that causes inflammation in certain parts of the spine. It usually starts in the sacroiliac joints, where the bottom of the spine joins the pelvis. Over time, the inflammation can cause the bones in the spine to fuse together, making the spine less flexible.

Here's a breakdown of the term:

Ankylosing: Means stiff or rigid.

Spondylitis: Means inflammation of the spine.

AS is a chronic condition that can cause significant pain and disability. However, with proper treatment, many people with AS can manage their symptoms and live a normal life.

Ankylosing spondylitis causes

The exact cause of ankylosing spondylitis is unknown. However, genetic factors, particularly the presence of the HLA-B27 gene, are strongly linked to the condition. This gene is found in most people with ankylosing spondylitis, but not everyone with the gene develops the condition.

While genetics play a significant role, other factors may also contribute to the development of ankylosing spondylitis, such as:

Immune system abnormalities: Some researchers believe that an overactive immune system may play a role.

Environmental factors: While not proven, some studies suggest that certain environmental factors may trigger the condition in people with a genetic predisposition.

It's important to note that further research is needed to fully understand the causes of ankylosing spondylitis.

Ankylosing spondylitis diagnosis

Diagnosing Ankylosing Spondylitis

Ankylosing spondylitis (AS) is a chronic inflammatory condition that primarily affects the spine. While there's no single definitive test, a combination of factors is often used to diagnose it.

Common Diagnostic Methods

Medical History: Your doctor will likely ask about:

Symptoms: Lower back pain, stiffness, fatigue, and limited range of motion.

Family history: AS often runs in families.

Other symptoms: Eye inflammation (uveitis), skin rashes (psoriasis), and digestive problems.

Physical Exam: Your doctor will assess your spine's flexibility and look for signs of inflammation.

Blood Tests:

C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR): These tests measure inflammation in the body.

HLA-B27 gene: While not a definitive diagnosis, the presence of this gene can increase the likelihood of AS.

Imaging Tests:

X-rays: Can show changes in the sacroiliac joints, where the spine connects to the pelvis.

MRI: Provides more detailed images of the spine and surrounding tissues.

When to See a Doctor

If you experience persistent lower back pain, especially in the morning, that improves with activity but worsens with rest, consider consulting a healthcare professional. Early diagnosis can help manage symptoms and prevent complications.

Note: While these are common diagnostic methods, the specific tests and procedures may vary depending on your individual circumstances. It's essential to discuss any concerns with your doctor.

Would you like to know more about the symptoms, causes, or treatment options for ankylosing spondylitis?

Ankylosing spondylitis icd10

The ICD-10 code for ankylosing spondylitis is M45.

This code falls under the category of "Dorsopathies" and is used to classify and diagnose ankylosing spondylitis, a chronic inflammatory disease that primarily affects the spine.

Ankylosing spondylosis +

Ankylosing Spondylitis: A Brief Overview

Ankylosing spondylitis (AS) is a chronic inflammatory disease that primarily affects the spine. It causes inflammation in the joints, leading to stiffness, pain, and eventually, the fusion of vertebrae.

Symptoms

Back pain and stiffness: Often begins in the lower back and worsens with rest.

Reduced range of motion: Difficulty turning the head or bending over.

Fatigue: Feeling tired even with enough sleep.

Eye inflammation: Uveitis, which can cause blurred vision or eye pain.

Skin problems: Psoriasis or eczema.

Causes

The exact cause of AS is unknown, but it is believed to be a genetic predisposition combined with environmental factors. The HLA-B27 gene is often found in people with AS, but not everyone with the gene develops the condition.

Treatment

While there's no cure for AS, treatment focuses on managing symptoms and improving quality of life. This may include:

Medications: Nonsteroidal anti-inflammatory drugs (NSAIDs), disease-modifying antirheumatic drugs (DMARDs), and biologic agents.

Physical therapy: Exercises to maintain flexibility and strength.

Occupational therapy: Strategies to help with daily activities.

Surgery: In severe cases, spinal fusion surgery may be necessary.

If you're experiencing symptoms of ankylosing spondylitis, it's important to consult with a healthcare professional for a proper diagnosis and treatment plan.

Would you like to know more about specific treatment options or the impact of AS on daily life?

Hla b27

Here's a comprehensive explanation of HLA-B27, incorporating visual aids:

What is HLA-B27?

Human Leukocyte Antigen (HLA): A group of proteins on the surface of cells that help the immune system recognize foreign substances.

HLA-B27: A specific HLA protein that is more common in people with certain autoimmune diseases.

Why is HLA-B27 important?

Risk factor for autoimmune diseases: People with HLA-B27 are more likely to develop:

Ankylosing spondylitis (AS): A chronic inflammatory disease that affects the spine and joints.

Reactive arthritis: A joint inflammation triggered by an infection.

Psoriatic arthritis: A joint inflammation associated with psoriasis.

Crohn's disease: An inflammatory bowel disease.

Uveitis: Inflammation of the eye.

Not a definitive diagnosis: Having HLA-B27 doesn't guarantee you'll develop an autoimmune disease, and not having it doesn't mean you can't.

Consult a doctor: If you have symptoms of an autoimmune disease, discuss your HLA-B27 status with a healthcare professional.

Monitor your health: Regular check-ups can help detect early signs of disease and start treatment if necessary.

Remember: HLA-B27 is just one factor in the development of autoimmune diseases. Other genetic and environmental factors also play a role.

Uveitis

Uveitis: A Brief Overview

Uveitis is a condition characterized by inflammation of the uvea, the middle layer of the eye. This layer is composed of the iris, ciliary body, and choroid. Uveitis can cause a range of symptoms, including pain, redness, blurred vision, and light sensitivity.

Types of Uveitis

Anterior uveitis: This is the most common type, affecting the front part of the uvea (the iris).

Intermediate uveitis: This involves the middle part of the uvea.

Posterior uveitis: This affects the back part of the uvea (the choroid).

Causes

The exact cause of uveitis is often unknown. However, it can be associated with various factors, including:

Infections: Viruses, bacteria, fungi, and parasites can all cause uveitis.

Autoimmune diseases: Conditions like rheumatoid arthritis, lupus, and Crohn's disease can increase the risk.

Systemic diseases: Certain systemic diseases, such as sarcoidosis and tuberculosis, can be associated with uveitis.

Trauma or injury: Eye injuries can sometimes lead to inflammation.

Medications: Certain medications can cause uveitis as a side effect.

Symptoms

Symptoms of uveitis can vary depending on the type and severity of the condition. Common symptoms include:

Eye pain

Redness

Blurred vision

Light sensitivity

Floaters (dark spots in the field of vision)

Decreased vision

Treatment

Treatment for uveitis depends on the underlying cause and severity of the condition. It may involve:

Eye drops: Corticosteroid eye drops can help reduce inflammation.

Oral medications: In some cases, oral corticosteroids or other medications may be necessary.

Injections: Steroid injections into the eye may be used to treat severe uveitis.

Surgery: In rare cases, surgery may be required to treat uveitis complications.

It's important to seek prompt medical attention if you experience any of these symptoms. Early diagnosis and treatment can help prevent vision loss.

Would you like to know more about a specific aspect of uveitis, such as its causes, symptoms, or treatment options?

Ankylosing spondylitis radiology

Ankylosing Spondylitis Radiology

Ankylosing spondylitis (AS) is a chronic inflammatory condition that primarily affects the spine and sacroiliac joints. It can lead to the fusion of these joints, resulting in a condition known as "bamboo spine." Radiology plays a crucial role in diagnosing and monitoring the progression of AS.

Common Imaging Techniques

X-rays:

Early stages: May show subtle changes in the sacroiliac joints, such as sacroiliitis.

Later stages: Can reveal the characteristic "bamboo spine" appearance due to the fusion of vertebral bodies.

Other findings: Bone erosion, osteopenia, and enthesitis (inflammation of the points where tendons and ligaments attach to bone).

MRI:

Sensitive: Can detect early changes in the sacroiliac joints, even before they're visible on X-rays.

Soft tissue evaluation: Useful for assessing inflammation, bone marrow edema, and nerve involvement.

CT:

Bone detail: Provides excellent detail of the bony structures, especially in advanced cases.

Fractures: Can be helpful in detecting fractures in patients with AS, particularly in the cervical spine.

Radiological Features of AS

Sacroiliitis: Inflammation and erosion of the sacroiliac joints, typically bilateral and symmetrical.

Syndesmophytes: Bony growths that form along the spine, connecting adjacent vertebral bodies.

Bamboo spine: A fusion of the vertebral bodies, creating a bamboo-like appearance on X-rays.

Osteopenia: Decreased bone density.

Enthesitis: Inflammation of the entheses, particularly at the insertion of tendons and ligaments.

Scoring Systems

Several radiological scoring systems are used to assess the severity of AS:

BASRI (Bath Ankylosing Spondylitis Radiology Index): Quantifies radiological changes in the spine and sacroiliac joints.

SASSS (Stoke Ankylosing Spondylitis Spinal Score): Evaluates spinal changes, including syndesmophytes and ankylosis.

Remember: The choice of imaging modality depends on the stage of the disease, the clinical presentation, and the specific information needed. A healthcare professional will determine the most appropriate imaging technique for a given patient.

Would you like to know more about the symptoms of ankylosing spondylitis or its treatment options?

Spondyloarthritis ++

Spondyloarthritis (SpA)

Spondyloarthritis is a general term for a group of inflammatory diseases that affect the spine and its surrounding tissues. When accompanied by inflammation, it is specifically referred to as spondyloarthritis. SpA is a group of inflammatory rheumatic diseases that can affect various joints, the tissues around the joints, and sometimes other organs.

Why is SpA Important to Know About?

Prevalence: Millions of people worldwide live with SpA.

Chronic nature: It generally has a long-term course.

Disability: If left untreated, it can significantly limit daily living activities.

Comorbidities: It can occur alongside other health conditions such as inflammatory bowel disease and psoriasis.

What are the Symptoms of SpA?

SpA symptoms can vary from person to person, but the most common ones include:

Back pain: Back pain that is worse in the morning and tends to improve with movement.

Sacroiliac joint pain: Deep pain in the hip and lower back area.

Joint swelling and pain: Swelling and pain in joints like the knees, ankles, and shoulders.

Eye inflammation: Eye inflammation called uveitis.

Skin lesions: Skin lesions associated with psoriasis.

Bowel problems: Symptoms of inflammatory bowel disease such as diarrhea and abdominal pain.

What Causes SpA?

The exact cause of SpA is not fully understood, but it is believed to involve a combination of genetic predisposition, environmental factors, and immune system disorders.

How is SpA Diagnosed?

To make a diagnosis, your doctor may:

Physical examination: Assess your joint movement, tenderness, and swelling.

Blood tests: Check for inflammation markers and genetic factors like HLA-B27.

Imaging: Use X-rays, MRI, and other imaging techniques to evaluate joint damage.

How is SpA Treated?

The goal of SpA treatment is to reduce pain, control inflammation, and prevent joint damage. Treatment options include:

Medication: Nonsteroidal anti-inflammatory drugs (NSAIDs) and biologic agents.

Physical therapy: Exercises and other physical therapy techniques to improve joint mobility and strengthen muscles.

Lifestyle modifications: A healthy diet, regular exercise, and adequate sleep.

Living with SpA

Although SpA is a chronic disease, with proper treatment and lifestyle changes, it is possible to live an active and healthy life.

Remember, this information is a general guide. Please consult your doctor for a detailed diagnosis and treatment plan.

Ankylosing spondylitis blood test

Blood Tests for Ankylosing Spondylitis

Ankylosing spondylitis is a chronic inflammatory condition that primarily affects the spine. While there's no definitive blood test to diagnose it, certain blood tests can provide valuable clues.

Here are the most common blood tests used to help diagnose ankylosing spondylitis:

C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR): These tests measure inflammation in the body. Elevated levels of CRP or ESR can indicate inflammation, which is often present in ankylosing spondylitis.

HLA-B27: Many people with ankylosing spondylitis have the HLA-B27 gene. However, it's important to note that having this gene doesn't guarantee a diagnosis, and not everyone with ankylosing spondylitis has it.

It's crucial to remember that blood tests alone cannot definitively diagnose ankylosing spondylitis. A diagnosis typically involves a combination of factors, including:

Symptoms: Persistent back pain, especially in the lower back, that is worse at night and improves with activity.

Physical examination: A doctor may assess joint stiffness, limited range of motion, and other physical signs.

Imaging tests: X-rays, MRI, or CT scans can help visualize changes in the spine, such as bone fusion.

If you're experiencing symptoms of ankylosing spondylitis, it's important to consult with a healthcare professional for a proper diagnosis and treatment plan.

Would you like to know more about the symptoms, treatment options, or other aspects of ankylosing spondylitis?

Perplexity

Ankylosing spondylitis pain (++)

Answer

Ankylosing spondylitis (AS) is a chronic inflammatory disease primarily affecting the spine and can also involve other joints. The condition is characterized by pain and stiffness, particularly in the lower back and hips. Understanding the nature of this pain, its symptoms, and treatment options is crucial for managing the condition effectively.

Nature of Pain in Ankylosing Spondylitis

The pain associated with ankylosing spondylitis is inflammatory in nature. It typically manifests as:

Lower back and hip pain: This pain often occurs in the early morning or after periods of inactivity, lasting for more than 30 minutes before easing with movement.

Nighttime discomfort: Many patients report that the pain can wake them during the night, necessitating movement to alleviate discomfort.

Stiffness: Morning stiffness is common, which improves with physical activity.

The pain can vary significantly among individuals, with some experiencing mild episodes while others endure chronic, severe pain. Inflammation may also affect peripheral joints, leading to additional symptoms such as tenderness in fingers or toes and discomfort in the ribs.

Symptoms Beyond Pain

In addition to pain and stiffness, ankylosing spondylitis can lead to a range of other symptoms:

Fatigue: Many individuals experience extreme tiredness.

Reduced flexibility: Over time, the condition can lead to a rigid spine due to the fusion of vertebrae.

Eye inflammation: Some patients may develop uveitis, causing vision changes and eye pain.

Other joint pain: Involvement of joints such as shoulders, knees, and ankles can occur.

Treatment Options

While there is currently no cure for ankylosing spondylitis, various treatment strategies aim to manage symptoms and improve quality of life:

Medications: Nonsteroidal anti-inflammatory drugs (NSAIDs) are commonly used to relieve pain and inflammation. For those who do not respond adequately to NSAIDs, biologic therapies targeting tumor necrosis factor (TNF) or interleukin-17 (IL-17) may be effective.

Physical therapy: Exercise is essential for maintaining mobility and posture. Patients are encouraged to engage in stretching and strengthening exercises tailored to their needs.

Lifestyle modifications: Regular physical activity can help alleviate symptoms. Patients are often advised against smoking, as it can exacerbate lung issues related to chest wall involvement.

Conclusion

Ankylosing spondylitis presents a complex challenge due to its inflammatory nature and potential for significant disability over time. Early diagnosis and a comprehensive treatment approach that includes medications, physical therapy, and lifestyle changes are vital for managing symptoms effectively and maintaining quality of life. Regular consultations with healthcare providers specializing in rheumatology are recommended for optimal management of this condition.

Ankylosing spondylitis symptoms (++)

Answer

Ankylosing spondylitis (AS) is a chronic inflammatory disease primarily affecting the spine and pelvis, leading to pain and stiffness. The symptoms can vary significantly among individuals but generally develop gradually over time.

Common Symptoms

1. Back Pain and Stiffness:

The most prominent symptom of AS is persistent pain and stiffness in the lower back and hips, particularly noticeable in the morning or after periods of inactivity. This discomfort often improves with physical activity but worsens with rest.

2. Joint Pain:

AS can cause inflammation in other joints, including the shoulders, knees, and feet. This may manifest as pain, swelling, tenderness, and warmth in the affected areas.

3. Enthesitis:

This refers to inflammation where tendons or ligaments attach to bones, commonly affecting areas like the Achilles tendon and the ribs where they connect to the breastbone. Symptoms may include localized pain and tenderness.

4. Fatigue:

Many individuals with AS experience significant fatigue, which can be debilitating and affect daily activities.

5. Eye Problems:

Uveitis, an inflammation of the eye, is a common complication associated with AS. Symptoms include eye pain, sensitivity to light, and blurred vision.

6. Other Symptoms:

Additional manifestations may include:

Chest pain and difficulty taking deep breaths if rib joints are involved.

Skin rashes, particularly psoriasis.

Abdominal pain or inflammatory bowel disease symptoms.

Progression of Symptoms

Symptoms often fluctuate, with periods of exacerbation (flares) followed by remission. While some individuals may only experience mild symptoms throughout their lives, others may face progressive stiffness and loss of spinal flexibility due to vertebrae fusion over time.

When to Seek Medical Attention

It is advisable to consult a healthcare provider if experiencing persistent lower back or buttock pain that worsens with inactivity or during the night but improves with exercise. Immediate attention is recommended for severe eye symptoms.

Understanding these symptoms can aid in early diagnosis and management of ankylosing spondylitis, which currently has no cure but can be managed effectively through various treatments.

Ankylosing spondylitis disease (++)

Overview of Ankylosing Spondylitis

Ankylosing spondylitis (AS) is a chronic inflammatory form of arthritis primarily affecting the spine and the sacroiliac joints, which connect the spine to the pelvis. The term "ankylosing" refers to the stiffness of the joints, while "spondylitis" denotes inflammation of the vertebrae. This condition can lead to severe pain and stiffness in the back, particularly in the lower regions, and may result in the fusion of vertebrae over time, leading to a rigid spine.

Symptoms

The symptoms of ankylosing spondylitis typically manifest gradually and can vary significantly among individuals. Common symptoms include:

Chronic back pain: Often worse during rest or at night and may improve with activity.

Stiffness: Particularly in the morning or after periods of inactivity.

Reduced spinal mobility: Difficulty bending or twisting due to stiffness.

Inflammation: Can also affect other joints such as hips, knees, and shoulders.

Systemic symptoms: These may include fatigue, weight loss, fever, and anemia.

In advanced cases, AS can lead to complications such as osteoporosis, spinal fractures, and involvement of other organs like the eyes (uveitis) and intestines (inflammatory bowel disease) .

Causes and Risk Factors

The exact cause of ankylosing spondylitis remains unclear; however, genetic factors play a significant role. A notable association exists with the HLA-B27 gene, which is present in a majority of individuals with AS but not exclusively responsible for its development. Environmental factors may also contribute to disease onset.

Risk factors include:

Age: Symptoms commonly appear between ages 17 and 35.

Gender: AS is more prevalent in men than women.

Family history: A family history of AS increases risk.

Diagnosis

Diagnosis typically involves a combination of clinical evaluation based on symptoms, medical imaging (such as X-rays or MRI), and blood tests to check for inflammatory markers and the presence of HLA-B27.

Treatment

While there is no cure for ankylosing spondylitis, various treatment options are available to manage symptoms and slow disease progression:

Medications:

Nonsteroidal anti-inflammatory drugs (NSAIDs) are commonly prescribed for pain relief.

Disease-modifying antirheumatic drugs (DMARDs) and biologic therapies targeting specific inflammatory pathways (e.g., TNF inhibitors) may be used for more severe cases.

Physical therapy: Tailored exercise programs can help maintain flexibility and posture.

Lifestyle modifications: Regular physical activity is encouraged to manage symptoms effectively.

Conclusion

Ankylosing spondylitis is a complex condition that requires ongoing management. Early diagnosis and a comprehensive treatment plan can significantly improve quality of life for those affected by this condition. Regular follow-up with healthcare providers is essential to monitor disease progression and adjust treatments as necessary.

What is Ankylosing spondylitis ++

Answer

Ankylosing spondylitis (AS) is a chronic inflammatory form of arthritis primarily affecting the spine and the sacroiliac joints, where the spine meets the pelvis. This condition can lead to significant stiffness and pain in the back, and in severe cases, it may result in the fusion of vertebrae, causing a rigid spine. The term "ankylosing" refers to the stiffening or fusion of bones, while "spondylitis" denotes inflammation of the vertebrae.

Causes and Risk Factors

The exact cause of ankylosing spondylitis remains unclear, but genetic factors play a significant role. A majority of individuals with AS possess a specific gene known as HLA-B27, which is associated with an increased risk of developing the condition. However, not everyone with this gene will develop AS, indicating that environmental factors may also contribute.

AS commonly manifests in young adults, typically between ages 17 and 35, and is more prevalent in men than women. Family history can increase susceptibility to the disease.

Symptoms

Symptoms of ankylosing spondylitis can vary widely among individuals but often include:

Chronic back pain: Typically worse at night or during periods of inactivity.

Stiffness: Especially noticeable in the morning or after sitting for long periods.

Reduced spinal flexibility: Leading to a stooped posture over time.

Pain in other joints: Such as hips and shoulders.

Fatigue and systemic symptoms: Including weight loss and fever.

In some cases, AS can also cause complications such as uveitis (eye inflammation), psoriasis (skin disease), and inflammatory bowel diseases like Crohn's disease.

Diagnosis and Treatment

Diagnosis of ankylosing spondylitis typically involves a combination of clinical evaluation, medical imaging (like X-rays or MRIs), and blood tests to identify inflammatory markers or the HLA-B27 antigen.

While there is no cure for AS, treatment options aim to relieve symptoms and improve quality of life. Common approaches include:

Medications: Nonsteroidal anti-inflammatory drugs (NSAIDs), corticosteroids, and biologics (such as TNF inhibitors) are often prescribed to manage pain and inflammation.

Physical therapy: Tailored exercises can help maintain spinal mobility and overall function.

Lifestyle modifications: Regular physical activity is encouraged to alleviate stiffness and improve posture.

Overall, while ankylosing spondylitis can significantly impact daily life, many individuals manage their symptoms effectively with appropriate treatment and lifestyle adjustments.

Ankylosing spondylitis treatment ++

Answer

Treatment of Ankylosing Spondylitis

The main goals of ankylosing spondylitis treatment are to:

Relieve symptoms like pain and stiffness

Maintain proper posture, flexibility, and strength

Halt or slow the progression of the disease

Prevent complications

Treatment typically involves a combination of:

Medications

Over-the-counter anti-inflammatory drugs (NSAIDs) to reduce inflammation and pain

Biologic medications that target specific immune responses and decrease inflammation

Janus kinase (JAK) inhibitors for disease unresponsive to other treatments

Corticosteroid injections to rapidly decrease inflammation

Physical Therapy

Exercises to improve flexibility, strength and posture

Physiotherapy using massage, manipulation and hydrotherapy

Lifestyle Measures

Maintaining good posture with assistive devices if needed

Stress management techniques like relaxation, yoga and tai chi

Quitting smoking, which can worsen symptoms

Maintaining a healthy weight to reduce joint stress

Surgery is rarely needed but may be considered for severely damaged joints or to correct spinal deformities.

While there is no cure for ankylosing spondylitis, early diagnosis and a combination of treatments can help manage symptoms, prevent complications, and allow many people to remain independent and active

As

Answer

The term "as" can refer to multiple contexts, including its use in language, sports, and specific organizations. Here are some notable interpretations:

Language Usage

Conjunction: In English, "as" is commonly used as a conjunction to indicate comparison or cause. For example, "as tall as" or "as a result."

Translation: In Turkish, "as" translates to various meanings depending on context, often relating to comparisons or conditions.

Sports

AS USA: This is a sports news outlet covering a variety of sports including NFL, NBA, soccer, and MLB. It provides breaking news, live events, and comprehensive sports information .

Diario AS: A prominent Spanish sports newspaper that focuses on football (soccer), basketball, and other sports news .

Organizations

AS Visa Solutions: A visa application center providing services for Schengen visa applications in countries like Hungary and Portugal .

AS Ecza Deposu: A pharmaceutical company based in Turkey .

As Medikal: A company specializing in pneumatic tube systems and medical supplies for dental services .

Each of these uses highlights the versatility of the term "as" across different domains.

As Ankylosing spondylitis +

Answer

Ankylosing spondylitis (AS) is a chronic inflammatory disease primarily affecting the axial skeleton, particularly the spine and sacroiliac joints. It is classified as a type of axial spondyloarthritis and is characterized by progressive spinal stiffness and chronic pain, which can lead to significant functional impairment over time.

Symptoms and Diagnosis

The hallmark symptoms of AS include:

Chronic Back Pain: Often begins in the lower back or gluteal region, typically worsening with rest and improving with physical activity.

Stiffness: Particularly noticeable in the morning or after periods of inactivity.

Postural Changes: As the disease progresses, individuals may develop a stooped posture due to spinal fusion.

Extra-Articular Symptoms: These can include eye inflammation (uveitis), fatigue, and gastrointestinal issues.

Diagnosis is based on clinical evaluation, medical imaging (such as X-rays or MRI), and laboratory tests that may show elevated inflammatory markers like C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR) .

Pathophysiology

The underlying mechanism of AS involves autoimmune processes, particularly related to the HLA-B27 antigen, which is present in a significant majority of patients. The disease is characterized by inflammation at the entheses (the sites where tendons and ligaments attach to bone), leading to pain and potential fusion of the spine over time. Key inflammatory cytokines involved include tumor necrosis factor-alpha (TNF-α) and interleukin-1 (IL-1) .

Treatment Options

While there is no cure for AS, various treatments aim to alleviate symptoms and slow disease progression:

Medications:

Non-Steroidal Anti-Inflammatory Drugs (NSAIDs): First-line treatment for managing pain and inflammation.

Disease-Modifying Anti-Rheumatic Drugs (DMARDs): Such as TNF inhibitors (e.g., etanercept, infliximab) that target specific inflammatory pathways.

Corticosteroids: Used for acute flares or severe symptoms.

Physical Therapy: Tailored exercise programs can improve flexibility, reduce stiffness, and enhance overall function. Hydrotherapy is also beneficial for pain management .

Surgery: In cases of severe joint damage or deformity, surgical interventions may be considered .

Prognosis

The progression of ankylosing spondylitis varies widely among individuals. Some may experience periods of remission with minimal symptoms, while others may face increasing disability due to spinal fusion and other complications. Although AS is associated with an increased risk of cardiovascular disease and osteoporosis, modern treatments have significantly improved outcomes and quality of life for many patients .

In summary, ankylosing spondylitis is a complex condition requiring a multifaceted approach to management, combining medication, physical therapy, and lifestyle adaptations to optimize patient outcomes.

Ankylosing spondylitis meaning ++

Answer

Ankylosing spondylitis (AS) is a chronic inflammatory type of arthritis primarily affecting the axial skeleton, particularly the spine and sacroiliac joints (where the spine meets the pelvis). The term "ankylosing" refers to stiffness or rigidity, "spondyl" pertains to the spine, and "itis" indicates inflammation. AS is characterized by long-term inflammation that can lead to pain, stiffness, and eventual fusion of the vertebrae, resulting in a rigid spine.

Symptoms

The hallmark symptoms of ankylosing spondylitis include:

Chronic back pain: Often worse at night and improves with activity.

Stiffness: Particularly noticeable in the morning or after periods of inactivity.

Postural changes: Such as a stooped posture due to spinal stiffness.

Peripheral joint involvement: Pain may also occur in other joints like hips and knees.

Extra-articular manifestations: These can include eye inflammation (uveitis), skin issues (like psoriasis), and gastrointestinal disorders (such as inflammatory bowel disease) .

Pathophysiology

The exact cause of AS remains unknown, but it is believed to involve genetic factors, particularly the presence of the HLA-B27 antigen, which is found in a significant proportion of individuals with the condition. The disease is thought to be autoimmune in nature, where the immune system mistakenly attacks the body's own tissues, leading to inflammation at sites where tendons and ligaments attach to bones (entheses) .

Diagnosis and Management

Diagnosis typically involves a combination of clinical evaluation, imaging studies (like X-rays or MRI), and laboratory tests. While there is no cure for AS, treatment focuses on managing symptoms and maintaining mobility. Common approaches include:

Medications: Nonsteroidal anti-inflammatory drugs (NSAIDs), disease-modifying antirheumatic drugs (DMARDs), and biologics like TNF inhibitors.

Physical therapy: To improve flexibility and posture.

Exercise: Regular physical activity is crucial for managing symptoms and preventing stiffness .

In summary, ankylosing spondylitis is a serious condition that can significantly impact quality of life but can be managed effectively with appropriate medical care and lifestyle adjustments.

Ankylosing spondylitis causes ++

Answer

Ankylosing spondylitis (AS) is a chronic inflammatory disease primarily affecting the spine and sacroiliac joints, leading to pain and stiffness. The exact cause of AS remains unknown, but several factors are believed to contribute to its development.

Genetic Factors

A significant association exists between AS and the HLA-B27 gene, a genetic variant found in a large proportion of individuals with the condition. Studies indicate that over 80% of people with AS carry this gene variant, particularly among white populations. However, having the HLA-B27 gene does not guarantee the development of AS; many individuals with this gene never experience the disease, suggesting that additional factors are involved.

Environmental Factors

While genetics play a crucial role, environmental factors are also thought to influence the onset of AS. Researchers have not pinpointed specific environmental triggers, but they suggest that infections or other immune responses might initiate or exacerbate the condition in genetically predisposed individuals.

Other Risk Factors

Several additional risk factors can increase susceptibility to AS:

Family History: A family history of ankylosing spondylitis or related autoimmune diseases increases the risk.

Age and Gender: AS typically manifests in young adults, particularly males, often before the age of 45.

Related Conditions: Individuals with other autoimmune diseases such as Crohn's disease, ulcerative colitis, or psoriasis may have a higher likelihood of developing AS.

Understanding these causes and risk factors is essential for early diagnosis and management of ankylosing spondylitis, as there is currently no cure for the disease. Treatment focuses on alleviating symptoms and maintaining mobility through medication, physical therapy, and lifestyle adjustments

Ankylosing spondylitis diagnosis ++

Answer

Ankylosing spondylitis (AS) is a chronic inflammatory disease primarily affecting the spine and sacroiliac joints, often leading to significant pain and stiffness. Diagnosing AS can be challenging due to its gradual onset and the absence of a single definitive test. The diagnostic process typically involves a combination of medical history, physical examination, imaging studies, and laboratory tests.

Initial Assessment

Medical History: The physician will inquire about the patient's symptoms, including:

Duration and location of pain.

Factors that exacerbate or alleviate the pain.

Family history of AS or related conditions.

Specific characteristics of the pain, such as whether it improves with exercise or worsens at night.

Physical Examination: The doctor will assess:

Joint flexibility and movement.

Signs of inflammation in the spine and other joints.

Chest expansion during deep breathing to check for rib stiffness.

Imaging Studies

Imaging plays a crucial role in diagnosing AS:

X-rays: These are used to identify changes in the sacroiliac joints. However, early-stage AS may not show visible changes on X-rays.

MRI: This imaging technique is more sensitive than X-rays and can detect early signs of inflammation in the sacroiliac joints before structural changes occur.

Laboratory Tests

While there is no single test for AS, several laboratory tests can provide supporting evidence:

HLA-B27 Gene Test: Many individuals with AS carry this gene, but its presence alone does not confirm the diagnosis since not everyone with the gene develops AS.

Inflammation Markers: Blood tests may check for elevated levels of inflammatory markers like C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR), which can indicate inflammation but are not specific to AS.

Diagnostic Criteria

A diagnosis of ankylosing spondylitis is often confirmed if:

An X-ray shows inflammation in the sacroiliac joints (sacroiliitis).

The patient exhibits at least one of the following:

Lower back pain lasting more than three months that improves with exercise but not with rest.

Limited movement in the lower back.

Reduced chest expansion compared to expected values for age and sex.

If imaging tests do not show clear signs of AS but symptoms persist, a diagnosis may still be considered based on clinical criteria, particularly if the HLA-B27 gene is present.

In conclusion, diagnosing ankylosing spondylitis requires a comprehensive approach that combines patient history, physical examination, imaging studies, and laboratory tests to effectively identify this complex condition. Early diagnosis is crucial for managing symptoms and preventing long-term complications associated with AS.

Ankylosing spondylitis icd10 ++

Answer

Ankylosing spondylitis is classified under the ICD-10 coding system with the primary code M45. This classification includes various subcategories that specify the affected regions of the spine. Below are the details of the ICD-10 codes related to ankylosing spondylitis:

Main ICD-10 Code

M45: Ankylosing spondylitis

Subcategories

M45.0: Ankylosing spondylitis of multiple sites in the spine

M45.1: Ankylosing spondylitis of occipito-atlanto-axial region

M45.2: Ankylosing spondylitis of cervical region

M45.3: Ankylosing spondylitis of cervicothoracic region

M45.4: Ankylosing spondylitis of thoracic region

M45.5: Ankylosing spondylitis of thoracolumbar region

M45.6: Ankylosing spondylitis of lumbar region

M45.7: Ankylosing spondylitis of lumbosacral region

M45.8: Ankylosing spondylitis sacral and sacrococcygeal region

M45.9: Ankylosing spondylitis of unspecified sites in spine

These codes are used for medical classification and billing purposes, helping healthcare providers accurately document and treat this condition

Ankylosing spondylosis++

Answer

Ankylosing spondylitis (AS) is a chronic inflammatory disease primarily affecting the spine and large joints, leading to pain and stiffness. It is classified as a type of axial spondyloarthritis and is characterized by inflammation at the sacroiliac joints, where the spine connects to the pelvis. Over time, AS can cause vertebrae to fuse, resulting in a rigid spine and potentially a hunched posture.

Symptoms

The symptoms of ankylosing spondylitis typically appear gradually, with a peak onset between the ages of 20 and 30. Common symptoms include:

Chronic back pain: Often worse at night or in the early morning, improving with movement.

Stiffness: Particularly in the lower back and hips, especially after periods of inactivity.

Reduced spinal mobility: Difficulty bending or twisting.

Fatigue: General tiredness that may accompany other symptoms.

Systemic symptoms: These may include weight loss, fever, and anemia.

Eye inflammation: Such as uveitis, which can cause pain and sensitivity to light.

Causes and Risk Factors

The exact cause of ankylosing spondylitis is not fully understood, but it is believed to involve genetic predisposition. A significant number of individuals with AS carry the HLA-B27 antigen, although not all who possess this gene will develop the condition. AS tends to affect more men than women and often arises in young adults aged 17 to 35.

Pathophysiology

Ankylosing spondylitis is considered an autoimmune or autoinflammatory disorder. The inflammation primarily affects the spine but can also involve other areas such as the hips, shoulders, and places where tendons attach to bones. In response to inflammation, new bone formation may occur, leading to fusion of vertebrae.

Diagnosis

Diagnosis typically involves a combination of patient history, physical examination, imaging studies (such as X-rays or MRI), and blood tests for inflammatory markers. The absence of rheumatoid factor distinguishes AS from other types of arthritis.

Treatment

While there is no cure for ankylosing spondylitis, treatment focuses on managing symptoms and slowing disease progression. Common treatment options include:

Nonsteroidal anti-inflammatory drugs (NSAIDs): To relieve pain and reduce inflammation.

Disease-modifying antirheumatic drugs (DMARDs): Such as TNF inhibitors that target specific pathways involved in inflammation.

Physical therapy: To maintain mobility and improve posture.

Exercise: Regular physical activity is encouraged to help manage stiffness and pain.

Complications

Over time, ankylosing spondylitis can lead to complications such as:

Spinal fusion: Resulting in a loss of flexibility.

Osteoporosis: Increasing the risk of fractures.

Heart problems: Inflammation can affect heart valves and increase cardiovascular risk.

Eye issues: Including uveitis or other inflammatory eye conditions.

In summary, ankylosing spondylitis is a complex condition that requires ongoing management to alleviate symptoms and maintain quality of life. Early diagnosis and tailored treatment plans are crucial for effective management of this chronic disease.

Hla b27 ++

HLA-B27 Overview

HLA-B27 is a genetic marker that belongs to the human leukocyte antigen (HLA) system, which plays a crucial role in the immune system. It is particularly significant due to its association with ankylosing spondylitis (AS) and other spondyloarthropathies. The HLA-B27 gene is located on the short arm of chromosome 6 and encodes a protein that is involved in the regulation of the immune response.

HLA-B27 and Ankylosing Spondylitis Relationship

What is Ankylosing Spondylitis?

Ankylosing spondylitis is a chronic inflammatory disease primarily affecting the spine and sacroiliac joints. It leads to inflammation, pain, and eventually stiffness in these areas. Individuals with the HLA-B27 gene have a significantly higher risk of developing ankylosing spondylitis, although not everyone who carries the gene will develop the disease.

HLA-B27 Testing

The HLA-B27 test is a genetic test that determines the presence of this antigen in an individual’s blood. A positive result can provide important clues for diagnosing ankylosing spondylitis, but it is not definitive on its own. The test is usually conducted when a patient exhibits symptoms of AS. In Turkey, approximately 70-80% of patients with ankylosing spondylitis test positive for HLA-B27, while in some Eastern European countries, this percentage can rise to 95%.

Clinical Manifestations Associated with HLA-B27

In individuals who are HLA-B27 positive, there is often a higher incidence of extra-articular manifestations. For instance, conditions such as uveitis (inflammation of the eye) are more common in these individuals. There is also a correlation between HLA-B27 positivity and inflammatory back pain; this type of pain tends to be more prevalent in HLA-B27 positive individuals compared to those who are negative.

Genetic Risks

The prevalence of HLA-B27 in the population correlates with the incidence of ankylosing spondylitis. In healthy populations, the prevalence of HLA-B27 positivity ranges from 7-10%, while the occurrence of ankylosing spondylitis is about 0.5%. Among first-degree relatives of individuals who are HLA-B27 positive, the risk of developing ankylosing spondylitis can increase to 15-20%, whereas it remains at 1-5% for those who are negative.

In conclusion, while the HLA-B27 test is an important tool in diagnosing ankylosing spondylitis, it should be interpreted alongside other clinical findings for an accurate diagnosis.

Uveitis ++

Uveitis is a significant condition characterized by inflammation of the uvea, which is the middle layer of the eye. This layer consists of three main parts: the iris, ciliary body, and choroid. Uveitis can affect one or more of these areas and is categorized into four main types based on the specific location of inflammation: anterior, intermediate, posterior, and panuveitis (involving all parts) .

Symptoms

The symptoms of uveitis can vary depending on the affected area but commonly include:

Eye redness

Pain in and around the eye

Blurry vision

Sensitivity to light (photophobia)

Floaters (dark spots in vision)

Symptoms may develop suddenly and can worsen quickly, making prompt medical attention crucial to prevent complications such as permanent vision loss .

Causes

Uveitis can arise from various causes, including:

Infections (e.g., herpes, syphilis, tuberculosis)

Autoimmune diseases (e.g., rheumatoid arthritis, sarcoidosis)

Injury to the eye

Idiopathic origins, where no specific cause is identified .

Approximately 50% to 70% of uveitis cases are classified as idiopathic, meaning their exact cause remains unknown .

Diagnosis and Treatment

Diagnosis typically involves a comprehensive eye examination by an ophthalmologist, who will assess symptoms, medical history, and possibly perform additional tests to identify underlying causes. Treatment strategies often include:

Corticosteroids (topical or injected) to reduce inflammation

Immunosuppressive drugs for chronic cases

Addressing any underlying conditions contributing to uveitis .

Early intervention is critical; untreated uveitis can lead to severe complications such as glaucoma, cataracts, retinal detachment, and permanent vision loss .

In summary, uveitis is a serious eye condition requiring immediate medical attention. Understanding its symptoms, potential causes, and treatment options can significantly influence outcomes for those affected.

Ankylosing spondylitis radiology ++

Imaging in Ankylosing Spondylitis

Imaging plays a crucial role in the diagnosis and management of ankylosing spondylitis (AS). The characteristic radiographic and magnetic resonance imaging (MRI) findings are essential for establishing the diagnosis and monitoring disease progression. The main imaging modalities used in AS are:

Radiography

Radiographs are the conventional method for assessing structural damage in the spine and sacroiliac joints.

Early radiographic findings include erosions, sclerosis, and widening of the sacroiliac joints, with changes typically starting on the iliac side.

As the disease progresses, syndesmophytes (vertical ossification along the sides of the vertebral bodies) and ankylosis of the sacroiliac and spinal joints occur, leading to the classic "bamboo spine" appearance.

MRI

MRI is more sensitive than radiography in detecting early inflammatory changes, such as bone marrow edema, synovitis, and enthesitis.

On MRI, active inflammation is characterized by subchondral bone marrow edema, which appears as low signal intensity on T1-weighted images and high signal on STIR/T2-weighted images.

MRI can detect inflammatory lesions in the sacroiliac joints and spine, including sacroiliitis, spondylitis, and spondylodiscitis, even before structural changes are visible on radiographs.

MRI is also useful for monitoring disease activity and response to treatment.

CT

CT is more sensitive than radiography for detecting chronic structural changes, such as erosions, sclerosis, and ankylosis of the sacroiliac joints.

CT is the imaging modality of choice for evaluating suspected spinal fractures in patients with advanced AS.

Other imaging modalities, such as ultrasonography and scintigraphy, have limited roles in the assessment of axial involvement in AS.

In summary, radiography remains the standard method for assessing structural damage, while MRI is crucial for detecting early inflammatory changes and monitoring disease activity. CT has a complementary role in evaluating chronic structural changes and suspected spinal fractures

Spondyloarthritis ++

Answer

Spondyloarthritis (SpA) is a term encompassing a group of inflammatory rheumatic diseases that primarily affect the spine and other joints. It is characterized by joint inflammation, particularly in the axial skeleton, leading to symptoms such as back pain and stiffness. The condition is often associated with the HLA-B27 gene, although not all individuals with this gene will develop the disease.

Types of Spondyloarthritis

Spondyloarthritis includes several subtypes, each with distinct features:

Ankylosing Spondylitis (AS): The most recognized form, AS primarily affects the spine and sacroiliac joints, leading to potential spinal fusion.

Axial Spondyloarthritis: This term encompasses AS and non-radiographic axial spondyloarthritis (nr-axSpA), where joint damage is not visible on X-rays.

Peripheral Spondyloarthritis: Involves inflammation of joints outside the spine, such as in the arms and legs.

Psoriatic Arthritis (PsA): Often associated with psoriasis, PsA can affect both peripheral joints and the spine.

Reactive Arthritis: Triggered by infections, this type may resolve on its own after the underlying infection is treated.

Enteropathic Arthritis: Linked to inflammatory bowel diseases like Crohn's disease and ulcerative colitis, affecting both the spine and peripheral joints.

Undifferentiated Spondyloarthritis: Symptoms align with spondyloarthritis but do not fit specific subtypes.

Symptoms

The hallmark symptom of spondyloarthritis is inflammatory back pain, which typically:

Starts gradually and persists for at least three months.

Improves with physical activity but worsens with rest.

Often includes morning stiffness that eases throughout the day.

Other symptoms may include:

Pain and swelling in peripheral joints.

Enthesitis (inflammation at tendon or ligament attachment sites).

Fatigue and general discomfort.

Diagnosis

Diagnosing spondyloarthritis involves a combination of clinical evaluation and imaging studies. Key diagnostic criteria include:

Presence of inflammatory back pain.

Limited spinal mobility.

Radiological evidence of sacroiliitis (inflammation of the sacroiliac joints) on X-rays or MRI scans.

Blood tests for HLA-B27 can support diagnosis but are not definitive since many patients may be HLA-B27 negative.

Treatment

Management of spondyloarthritis typically includes:

Physical Therapy: To improve mobility and reduce stiffness.

Medications:

Nonsteroidal anti-inflammatory drugs (NSAIDs) are first-line treatments for pain relief.

Disease-modifying antirheumatic drugs (DMARDs) may be prescribed if NSAIDs are ineffective.

Biologics targeting specific inflammatory pathways can be effective for severe cases.

Surgical interventions, such as hip replacement or spinal surgery, are rarely needed but may be considered in severe cases.

Prognosis

With appropriate treatment, many individuals with spondyloarthritis can lead normal, productive lives. Regular exercise and lifestyle modifications, including smoking cessation, are recommended to manage symptoms effectively. While some patients may experience significant progression leading to complications like spinal fusion, others may have milder forms that remain manageable

Ankylosing spondylitis blood test ++

Ankylosing Spondylitis Blood Tests

There are several blood tests that can help diagnose ankylosing spondylitis (AS), but no single test can definitively confirm the condition. Doctors use a combination of blood tests, imaging scans, and a physical exam to diagnose AS.

HLA-B27 Genetic Test

The HLA-B27 blood test checks for a specific gene variant that is found in 90-95% of people with AS. However, having the HLA-B27 gene does not necessarily mean you have AS, as about 7% of the general population also carries this gene. Additionally, some people with AS do not have the HLA-B27 gene.

Inflammation Blood Tests

Other blood tests look for markers of inflammation in the body, which can be a sign of AS:

C-reactive protein (CRP) test measures levels of CRP, a protein produced by the liver that increases with inflammation. CRP is often elevated in AS.

Erythrocyte sedimentation rate (ESR or sed rate) measures how quickly red blood cells fall to the bottom of a test tube. A faster than normal rate suggests inflammation.

Plasma viscosity (PV) test measures the thickness of the blood, which increases with inflammation. A positive PV result could indicate AS or another inflammatory condition.

However, these inflammation tests are not specific to AS. Many other conditions can also cause elevated CRP, ESR, and PV levels.

In summary, while blood tests can provide supportive evidence for an AS diagnosis, they cannot diagnose the condition on their own. Your doctor will need to consider your symptoms, physical exam findings, and imaging test results along with your blood work to determine if you have AS